Condoms in the prevention of HIV and AIDS

Lucian Jayasuriya¹

Within a quarter of a century HIV/AIDS has devastated mankind.

It is the most critical health issue facing mankind today. AIDS is the leading cause of death in Africa and the fourth leading cause of death worldwide. Millions of children have been orphaned especially in Africa.

What are the strategies available to control this scourge?

There is no vaccine in sight. Treatment with highly active anti retroviral therapy only prolongs life and prevents mother to child transmission to some extent..

Prevention remains the answer to the problem.

Theoretically we say prevention is by ABC – abstinence, be faithful and condoms. We mouth the first two and depend on condom promotion to prevent HIV transmission. This strategy has obviously failed because the numbers getting infected are increasing continuously. Sadly however, those who lead the campaign to control HIV/AIDS internationally do not seem to want to face reality. They seem to want to have the cake and eat it.

They want to promote sexual activity as well as control the spread of sexually transmitted infections.

To prevent transmission of HIV by sexual intercourse condoms of good quality have to be used correctly and consistently. This is the crunch. It is obvious that these criteria of usage are often not fulfilled. The end result is that the condom usage results in acquiring infection sooner or later. Therefore condom promotion should not be a strategy for prevention of HIV transmission.

The pandemic in Africa is due to the fact that sex before marriage and sex after marriage are considered as the norm in these cultures.

Sri Lankan culture does not consider pre- marital and extra marital sex as the norm. This may a reason that Sri Lanka remains a low prevalence country for HIV/AIDS.

Abstinence before marriage and being faithful in marriage should be promoted to keep our low prevalence status. Usage of condoms should not be promoted except to commercial sex workers and men who have sex with men.

¹Founder President, College of Venereologists of Sri Lanka

The first intravenous drug user with HIV infection in Sri Lanka

MPW Gunathilake¹, A Wijewickrama², S Mananwatte³, S Samarakoon⁴

Introduction

Blood to blood transmission of HIV either by needle sharing intravenous drug use or blood transfusion is much more effective than sexual transmission.

Explosive epidemics of HIV, related to intravenous drug use have been observed in many parts of the world.

Injecting drug use is not reported as a problem in Sri Lanka, at present. The National Dangerous Drugs Control Board estimates 40,000 heroin users in the country and

200,000 cannabis users, only 2% are estimated to be injecting drug users. While the World Bank estimates 7.5% of the illicit drug users are injectors.¹

We are reporting the first intravenous drug user with HIV infection in this country.

Case Report

In July 2004, HIV-2 infection was detected in a 36 year old Middle East returnee. He admitted to have been an intravenous drug user since 1992, from the time he was

¹Senior Register in Venereology, Central STD Clinic National STD/AIDS Control Programme, ² Consultant Physician, Infectious diseases Hospital, Gothatuwa, ³ Consultant Microbiologist, ⁴Consultant Venereologist, Central STD Clinic, National STD/AIDS Control Programme.

imprisoned in a Middle Eastern country, during which period he claimed to have been introduced to IV drugs by other prison inmates.

He denied any risky sexual behaviors while in or out of prison.

In late January 2004, he developed night sweats, fever and difficulty in swallowing, suggestive of primary HIV infection. However HIV antibodies were negative at that time.

He returned to Sri Lanka in July 2004 and tested positive for HIV infection at a private hospital.

The Line Blot assay done for confirmation of HIV infection at the NSACP, showed the typical gp 105 and gp 36 bands, which denotes HIV-2 infection.

He was successfully treated for oral candidiasis after which he was asymptomatic but the CD4 count was 78 cells/ μ l, rendering him a candidate for highly active anti retroviral therapy (HAART)

He was commenced on HAART (zidovudine, lamivudine, and indinavir boosted with ritonavir) with cotrimoxazole prophylaxis.

His adherence to treatment was good and at six months, his CD4 count was observed to be 228 cells/ μ l. His response to treatment continues to be very good with a CD4 count of 721/ μ l in January 2007.

¹Senior Registrar in Venereology, National STD/AIDS Control Programme, ²Consultant Physician, Infectious Diseases Hospital, ³Consultant Microbiologist, National STD/AIDS Control Programme, ⁴Consultant Venereologist, National STD/AIDS Control Programme

Discussion

In 1986, three years after the discovery of the human immunodeficiency virus, a second type of HIV, called HIV-2 was isolated from AIDS patients in West Africa².

The prevalence of HIV-2 remains highest in West Africa, and in countries that have socio-economic ties with the area.² Although both HIV-1 and HIV-2 belong to the lentivirus subfamily, and their genomic organization is similar ³, HIV-2 is only about 40% similar to HIV-1 in nucleotide sequence and is more closely related to Simian immunodeficiency virus with approximately 75% nucleotide sequence similarity⁴.

The clinical and biological characteristics of infection with these two viruses show substantial differences as well as similarities⁵.

Like HIV-1, HIV-2 has been shown to use CD4 cells as a primary receptor and several chemokine receptors as co-receptors, for infection of target cells⁶. HIV-1 uses either CCR5 or CXCR4 as major co-receptors, but can also use other molecules to a lesser extent, such as CCR 3. Interestingly several HIV-2 strains have been shown to infect certain CD4 negative cell lines, most by using CXCR4 or CCR5 alone and are better at doing so than HIV-1⁷.

Transmission of HIV-2 occurs mainly by heterosexual contact or contact with infected blood. Efficiency of heterosexual transmission of HIV-2 is estimated to be five times lower than that of HIV-1⁸. Additionally the rate of viral shedding in cervical samples was shown to be 16% in HIV-2 positive women, compared with 36.4% in HIV-1 positive women⁹. Transmission from mother to child during pregnancy or through breast milk is much less likely with HIV-2. Although vertical transmission occurs in approximately one third of untreated HIV-1 positive pregnancies¹⁰, most studies have shown that vertical transmission of HIV-2 occurs rarely. (About 0-4% of all HIV-2 positive pregnancies.)

HIV-2 causes AIDS but the period between infection and development of AIDS appears to be substantially longer in than that for HIV-1¹¹. This difference in disease progression could, in part, be due to lower viral loads in those infected with HIV-2 than in those infected with HIV-1. However, once disease develops, patients with HIV-2 generally have most of the signs and symptoms of HIV-1 patients.

Presenting symptoms of HIV-2 infection include generalized lymphadenopathy, candidiasis, and pulmonary disease, generalized pruritic dermatitis, weight loss of greater than 10% body weight, and diarrhoea or fever for greater than one month¹². Those infected with HIV-2, sometimes complain of fever, diarrhea, and/or oral candidiasis around the time of sero conversion¹².

Not all the drugs used to treat HIV-1 infection are as effective against HIV-2. Non nucleoside reverse transcriptase inhibitors (NNRTIs) are largely ineffective against HIV-2¹³. Nucleoside reverse transcriptase inhibitors (NRTIs) are effective but may be less so than in HIV-1¹³. Protease Inhibitors appear to be effective generally¹³.

Monitoring the treatment response of patients infected with HIV-2 is difficult. No FDA- licensed HIV-2 viral load assay is available yet.

Hence monitoring of disease progression and therapy, in the absence of viral load assay, are clinical assessment and CD4 count.

The present knowledge on HIV-2 is limited due to small number of patients with this infection worldwide.

References:

1. National Dangerous Drugs Control Board (NDDCB) 2005
2. Guyader CFM, Guetard D, Montagnier SM et al. Cloning and polymorphism of the human immunodeficiency virus type 2. *Nature* 1986; 324: 691-695
3. Guyader M, Emerman M, Sonigo P et al. Genome organization and transactivation of the human immunodeficiency virus type-2. *Nature* 1987;326:662-669.
4. Hirsch VM, Olmsted RA, Murphy-Corb M, et al An African primate lentivirus (SIVsm) closely related to HIV-2. *Nature* 1989;339:389-392
5. Kanki PJ, Biologic features of HIV-2. An update. *AIDS Clin Rev* 1991: vol 7-38
6. Rucker J, Edinger AL, Sharron M, et al. Utilization of chemokine receptors, orphan receptors and herpesvirus-encoded receptors by diverse human and simian immunodeficiency viruses. *J Virol* 1997;71:8999-9007.
7. Reeves JD, Hibbitts S, Simmons G et al. Primary HIV-2 isolates infect CD4 negative cells via CCR5 and CXCR4, *Journal of Virology* 1999;73; 9:7795-7804
8. Donnelly C, Leisenring W, Kanki P et al Comparison of transmission rates of HIV-1 and HIV-2 in a cohort of prostitutes in Senegal. *Bull Math Biol* 1993; 55:731-743.
9. Sankale JL, Mboup S, Essex ME et al. Genetic characterization of viral quasispecies in blood and cervical secretions of HIV-1- and HIV-2-infected women. *AIDS Res Hum Retroviruses* 1998; 14:1473-1481.
10. Cavaco-Silva P, Taveira NC, Lourenco MH et al. Vertical transmission of HIV-2. *Lancet* 1997; 349:177-178.
11. Ancelle R, Blettry O, Baglin AC et al. Long incubation period for HIV-2 infection. *Lancet* 1987; 1:688-689.
12. Norrgren H, Da Silva ZJ, Andersson S, et al. Clinical features, immunological changes and mortality in a cohort of HIV-2-infected individuals in Bissau, Guinea-Bissau. *Scand J Infect Dis* 1998; 30:323-329.
13. Schutten M, van der Ende ME, Osterhaus AD. Antiretroviral therapy in patients with dual infection with human immunodeficiency virus types 1 and 2. *N Engl J Med* 2000; 342:1758-1760